arXiv:1411.5257v1 [math.CA] 19 Nov 2014

An extension of a series containing Laguerre polynomials

A. K. Rathie[a] and R. B. Paris[b]

[a] Department of Mathematics, Central University of Kerala, Kasaragad 671123, Kerala, India
E-Mail: akrathie@cukerala.edu.in
[b] School of Engineering, Computing and Applied Mathematics, University of Abertay Dundee, Dundee DD1 1HG, UK
E-Mail: r.paris@abertay.ac.uk

Abstract

Expressions for the summation of the series involving the Laguerre polynomials

$$S_m(\pm\nu,\pm p)\equiv e^{-x}\sum_{n=0}^{\infty}\frac{x^n\,L_n^{(\nu)}(x)}{(1\pm\nu\pm p)_n}\frac{(f+m)_n}{(f)_n}$$

for any non-negative integers m and p are obtained in terms of generalized hypergeometric functions. These results extend previous work in the literature.

Mathematics Subject Classification: 33C15, 33C20

Keywords: Laguerre polynomials, generalized hypergeometric functions, generalized Kummer summation theorem

1. Introduction

The generalized Laguerre polynomials $L_n^{(\nu)}(x)$ are encountered in many branches of pure and applied mathematics. They form an orthogonal set on $[0,\infty)$ with the weight function $x^\nu e^{-x}$, and can be represented as a terminating confluent hypergeometric function ${}_1F_1$ in the form

$$L_n^{(\nu)}(x)=\frac{(\nu+1)_n}{n!}\,{}_1F_1(-n;\nu+1;x),$$

where $(a)_n$ denotes the Pochhammer symbol, or rising factorial, defined by $(a)_n=\Gamma(a+n)/\Gamma(a)$. The first three polynomials are given by

$$L_0^{(\nu)}(x)=1,\quad L_1^{(\nu)}(x)=1-x+\nu,\quad L_2^{(\nu)}(x)=\tfrac{1}{2}x^2-(\nu+2)x+\tfrac{1}{2}(\nu+1)(\nu+2).$$

In [3], Kim *et al.* obtained summation formulas for the series involving the generalized Laguerre polynomial $L_n^{(\nu)}(x)$ given by

$$\sum_{n=0}^{\infty}\frac{x^n\,L_n^{(\nu)}(x)}{(1\pm\nu\pm p)_n}$$

for integer p, where $0 \le p \le 5$. The case $p = 0$ had been considered originally by Exton [2]. Recently, Brychkov [1] has extended these results for any integer $p \ge 0$. Alternative expressions for $S(\pm\nu, \pm p)$, also for arbitrary integer $p \ge 0$, in terms of generalized hypergeometric functions of argument $-x^2$ have been obtained in [4]. These results differ from, and in some cases are simpler than, those obtained in [1]. The aim of this note is to extend the above evaluations to the case when there is a pair of numeratorial and denominatorial parameters differing by a positive integer m. We derive expressions for the summation of the series given by

$$S_m(\pm\nu, \pm p) \equiv e^{-x} \sum_{n=0}^{\infty} \frac{x^n\, L_n^{(\nu)}(x)}{(1 \pm \nu \pm p)_n} \frac{(f+m)_n}{(f)_n}$$

for any non-negative integer p and complex parameter $f \neq 0, -1, -2, \ldots$ when it is supposed that $1 \pm \nu \pm p \neq 0, -1, -2, \ldots$.

2. The case $m = 0$

We first summarize the results obtained in [3] in the case $m = 0$, where

$$S_0(\pm\nu, \pm p) = e^{-x} \sum_{n=0}^{\infty} \frac{x^n\, L_n^{(\nu)}(x)}{(1 \pm \nu \pm p)_n}. \tag{2.1}$$

Then, for $p = 0, 1, 2, \ldots$, we have

$$S_0(\nu, p) = \frac{(-)^p 2^{2\nu+p} \Gamma(1+\nu)}{\Gamma(1+2\nu+p)} \sum_{s=0}^{p} (-)^s \binom{p}{s} \left\{ A_0(\nu, p) F_0^{(1)} - \frac{4x(1+\nu) B_0(\nu, p)}{(1+\nu+p)(1+2\nu+p)} F_0^{(2)} \right\} \tag{2.2}$$

and

$$S_0(\nu, -p) = \frac{2^{2\nu-p} \Gamma(1+\nu-p)}{\Gamma(1+2\nu-p)} \sum_{s=0}^{p} \binom{p}{s} \left\{ A_0(\nu, -p) F_0^{(3)} - \frac{4x B_0(\nu, -p)}{1+2\nu-p} F_0^{(4)} \right\}, \tag{2.3}$$

where

$$A_0(\nu, p) = \frac{\Gamma(\nu + \frac{1}{2}p + \frac{1}{2}s + \frac{1}{2})}{\Gamma(\frac{1}{2} - \frac{1}{2}|p| + \frac{1}{2}s)}, \qquad B_0(\nu, p) = \frac{\Gamma(\nu + \frac{1}{2}p + \frac{1}{2}s + 1)}{\Gamma(\frac{1}{2}s - \frac{1}{2}|p|)},$$

and

$$F_0^{(1)} = {}_4F_5 \left[\begin{matrix} \frac{1}{2}+\frac{1}{2}\nu,\ 1+\frac{1}{2}\nu,\ \frac{1}{2}+\nu+\frac{1}{2}p+\frac{1}{2}s,\ \frac{1}{2}+\frac{1}{2}p-\frac{1}{2}s \\ \frac{1}{2},\ \frac{1}{2}+\frac{1}{2}\nu+\frac{1}{2}p,\ 1+\frac{1}{2}\nu+\frac{1}{2}p,\ \frac{1}{2}+\nu+\frac{1}{2}p,\ 1+\nu+\frac{1}{2}p \end{matrix} ; -x^2 \right],$$

$$F_0^{(2)} = {}_4F_5 \left[\begin{matrix} 1+\frac{1}{2}\nu,\ \frac{3}{2}+\frac{1}{2}\nu,\ 1+\nu+\frac{1}{2}p+\frac{1}{2}s,\ 1+\frac{1}{2}p-\frac{1}{2}s \\ \frac{3}{2},\ 1+\frac{1}{2}\nu+\frac{1}{2}p,\ \frac{3}{2}+\frac{1}{2}\nu+\frac{1}{2}p,\ 1+\nu+\frac{1}{2}p,\ \frac{3}{2}+\nu+\frac{1}{2}p \end{matrix} ; -x^2 \right],$$

$$F_0^{(3)} = {}_2F_3 \left[\begin{matrix} \frac{1}{2}+\nu-\frac{1}{2}p+\frac{1}{2}s,\ \frac{1}{2}+\frac{1}{2}p-\frac{1}{2}s \\ \frac{1}{2},\ \frac{1}{2}+\nu-\frac{1}{2}p,\ 1+\nu-\frac{1}{2}p \end{matrix} ; -x^2 \right], \quad F_0^{(4)} = {}_2F_3 \left[\begin{matrix} 1+\nu-\frac{1}{2}p+\frac{1}{2}s,\ 1+\frac{1}{2}p-\frac{1}{2}s \\ \frac{3}{2}, 1+\nu-\frac{1}{2}p,\ \frac{3}{2}+\nu-\frac{1}{2}p \end{matrix} ; -x^2 \right].$$

The function ${}_rF_{r+1}$ denotes the generalized hypergeometric function with r numeratorial and $r+1$ denominatorial parameters defined by

$${}_rF_{r+1} \left[\begin{matrix} a_1, \ldots, a_r \\ b_1, \ldots, b_{r+1} \end{matrix} ; z \right] = \sum_{k=0}^{\infty} \frac{(a_1)_n \ldots (a_r)_n}{(b_1)_n \ldots (b_{r+1})_n} \frac{z^n}{n!} \qquad (|z| < \infty).$$

When the sign of the parameter ν in the denominator of (2.1) is reversed, we have for $p = 0, 1, 2, \ldots$

$$S_0(-\nu,p) = \frac{(-2)^p}{p!}\sum_{s=0}^{p}(-)^r\binom{p}{s}\left\{C_0(-\nu,p)F_0^{(5)} - \frac{4xD_0(-\nu,p)}{(1+p)(1-\nu+p)}F_0^{(6)}\right\} \tag{2.4}$$

and

$$S_0(-\nu,-p) = 2^{-p}\sum_{s=0}^{p}\binom{p}{s}\left\{C_0(-\nu,-p)F_0^{(7)} - \frac{4xD_0(-\nu,-p)}{1-\nu-p}F_0^{(8)}\right\}, \tag{2.5}$$

where

$$C_0(\nu,p) = \frac{\Gamma(\frac{1}{2}\nu+\frac{1}{2}p+\frac{1}{2}s+\frac{1}{2})}{\Gamma(\frac{1}{2}\nu-\frac{1}{2}|p|+\frac{1}{2}s+\frac{1}{2})}, \qquad D_0(\nu,p) = \frac{\Gamma(\frac{1}{2}\nu+\frac{1}{2}p+\frac{1}{2}s+1)}{\Gamma(\frac{1}{2}\nu-\frac{1}{2}|p|+\frac{1}{2}s)},$$

and the functions $F_0^{(j)}$ $(5 \le j \le 8)$ are defined by

$$F_0^{(5)} = {}_3F_4\left[\begin{matrix}1,\ \frac{1}{2}+\frac{1}{2}\nu+\frac{1}{2}p-\frac{1}{2}s,\ \frac{1}{2}-\frac{1}{2}\nu+\frac{1}{2}p+\frac{1}{2}s\\ \frac{1}{2}+\frac{1}{2}p,\ 1+\frac{1}{2}p,\ \frac{1}{2}-\frac{1}{2}\nu+\frac{1}{2}p,\ 1-\frac{1}{2}\nu+\frac{1}{2}p\end{matrix};-x^2\right],$$

$$F_0^{(6)} = {}_3F_4\left[\begin{matrix}1,\ 1+\frac{1}{2}\nu+\frac{1}{2}p-\frac{1}{2}s,\ 1-\frac{1}{2}\nu+\frac{1}{2}p+\frac{1}{2}s\\ 1+\frac{1}{2}p,\ \frac{3}{2}+\frac{1}{2}p,\ 1-\frac{1}{2}\nu+\frac{1}{2}p,\ \frac{3}{2}-\frac{1}{2}\nu+\frac{1}{2}p\end{matrix};-x^2\right],$$

$$F_0^{(7)} = {}_2F_3\left[\begin{matrix}\frac{1}{2}-\frac{1}{2}\nu-\frac{1}{2}p+\frac{1}{2}s,\ \frac{1}{2}+\frac{1}{2}\nu+\frac{1}{2}p-\frac{1}{2}s\\ \frac{1}{2},\ \frac{1}{2}-\frac{1}{2}\nu-\frac{1}{2}p,\ 1-\frac{1}{2}\nu-\frac{1}{2}p\end{matrix};-x^2\right],$$

$$F_0^{(8)} = {}_2F_3\left[\begin{matrix}1-\frac{1}{2}\nu-\frac{1}{2}p+\frac{1}{2}s,\ 1+\frac{1}{2}\nu+\frac{1}{2}p-\frac{1}{2}s\\ \frac{3}{2},\ 1-\frac{1}{2}\nu-\frac{1}{2}p,\ \frac{3}{2}-\frac{1}{2}\nu-\frac{1}{2}p\end{matrix};-x^2\right].$$

We note that $C_0(-\nu,-p) = 1$ and $D_0(-\nu,-p) = \frac{1}{2}(s-p-\nu)$.

3. The case $m \neq 0$

We start with the transformation [3, (3.5)]

$$e^{-x}\sum_{n=0}^{\infty}\frac{(a_1)_n\ldots(a_P)_n}{(b_1)_n\ldots(b_Q)_n}(-xy)^n\,L_n^{(\nu)}(x) = \sum_{n=0}^{\infty}\frac{(-x)^n}{n!}\,{}_{P+2}F_Q\left[\begin{matrix}-n,-n-\nu,a_1,\ldots,a_P\\ b_1,\ldots,b_Q\end{matrix};y\right],$$

where P and Q are non-negative integers. In this, if we take $P = r$, $Q = r+1$, $y = -1$, and $a_1 = f_1+m_1,\ldots,a_r = f_r+m_r$, $b_1 = f_1,\ldots,b_r = f_r$ and $b_{r+1} = 1\pm\nu\pm p$, where m_s $(1 \le s \le r)$ are non-negative integers, then

$$e^{-x}\sum_{n=0}^{\infty}\frac{x^n\,L_n^{(\nu)}(x)}{(1\pm\nu\pm p)_n}\,\frac{(f_1+m_1)_n\ldots(f_r+m_r)_n}{(f_1)_n\ldots(f_r)_n}$$

$$= \sum_{n=0}^{\infty}\frac{(-x)^n}{n!}\,{}_{r+2}F_{r+1}\left[\begin{matrix}-n,-n-\nu,f_1+m_1,\ldots,f_r+m_r\\ 1\pm\nu\pm p,f_1,\ldots,f_r\end{matrix};-1\right]. \tag{3.1}$$

The ${}_{r+2}F_{r+1}$ series on the right-hand side of (3.1) can be evaluated with the help of the result given in (4.2) of [7]. However, to avoid overly complicated formulas, we restrict our attention here to the case of a single f-parameter ($r = 1$) and adopt a different procedure.

3.1 *A lemma connected with* $S_m(\pm\nu, \pm p)$

We consider the series

$$S_m(\pm\nu,\pm p) = e^{-x}\sum_{n=0}^{\infty}\frac{x^n\,L_n^{(\nu)}(x)}{(1\pm\nu\pm p)_n}\,\frac{(f+m)_n}{(f)_n} = \sum_{n=0}^{\infty}\frac{(-x)^n}{n!}\,{}_3F_2\left[\begin{matrix}-n,-n-\nu,f+m\\1\pm\nu\pm p,f\end{matrix};-1\right], \tag{3.2}$$

for positive integer m and non-negative integer p. Here, x is an arbitrary complex variable and f, ν are complex parameters such that $f \neq 0,-1,-2,\ldots$ and $1\pm\nu\pm p \neq 0,-1,-2,\ldots$.

To proceed we make use of the following lemma that expresses the ${}_3F_2$ series in (3.2) in terms of a finite Gauss hypergeometric series.

Lemma 1 *The following formula holds true*

$$S_m(\pm\nu,\pm p) = \sum_{r=0}^{m}\binom{m}{r}\frac{x^r}{(f)_r(1\pm\nu\pm p)_r}\sum_{n=0}^{\infty}\frac{(-x)^n}{n!}(n+\nu+1)_r\,{}_2F_1\left[\begin{matrix}-n,-n-\nu\\1\pm\nu\pm p+r\end{matrix};-1\right] \tag{3.3}$$

for non-negative integer m and p.

Proof. We write the ${}_3F_2(-1)$ series appearing on the right-hand side of (3.2) as a finite sum of ${}_2F_1(-1)$ series by means of the result [5, Eq. (7.2.3.15)]

$${}_3F_2\left[\begin{matrix}a,b,&f+m\\c,&f\end{matrix};x\right] = \sum_{r=0}^{m}\binom{m}{r}\frac{(a)_r(b)_r}{(c)_r(f)_r}\,x^r\,{}_2F_1\left[\begin{matrix}a+r,b+r\\c+r\end{matrix};x\right]$$

for positive integer m. Then it follows that

$${}_3F_2\left[\begin{matrix}-n,-n-\nu,f+m\\1\pm\nu\pm p,f\end{matrix};-1\right] = \sum_{r=0}^{m}(-)^r\binom{m}{r}\frac{(-n)_r(-n-\nu)_r}{(f)_r(1\pm\nu\pm p)_r}\,{}_2F_1\left[\begin{matrix}-n+r,-n-\nu+r\\1\pm\nu\pm p+r\end{matrix};-1\right]. \tag{3.4}$$

When $m > n$, we note that the sum on the right-hand side of (3.4) is over $0 \le r \le n$, since for $n+1 \le r \le m$ the factor $(-n)_r$ vanishes.

Substitution of (3.4) into (3.2) and use of $(-n)_r = (-)^r n(n-1)\ldots(n-r+1)$ then produces

$$\begin{aligned}
&S_m(\pm\nu,\pm p)\\
&= \sum_{r=0}^{m}\binom{m}{r}\frac{1}{(f)_r(1\pm\nu\pm p)_r}\sum_{n=r}^{\infty}\frac{(-x)^n}{(n-r)!}\,(-n-\nu)_r\,{}_2F_1\left[\begin{matrix}-n+r,-n-\nu+r\\1\pm\nu\pm p+r\end{matrix};-1\right]\\
&= \sum_{r=0}^{m}\binom{m}{r}\frac{(-x)^r}{(f)_r(1\pm\nu\pm p)_r}\sum_{n=0}^{\infty}\frac{(-x)^n}{n!}\,(-n-\nu-r)_r\,{}_2F_1\left[\begin{matrix}-n,-n-\nu\\1\pm\nu\pm p+r\end{matrix};-1\right].
\end{aligned}$$

Finally, employing the fact that $(-n-\nu-r)_r = (-)^r(n+\nu+1)_r$ we obtain the result stated in the lemma. □

The ${}_2F_1$ series on the right-hand side of (3.3) can now be evaluated with the help of the generalized Kummer summation theorem in the form [6]

$${}_2F_1\left[\begin{matrix}a,b\\1+a-b+j\end{matrix};-1\right] = 2^{-2b+j}\frac{\Gamma(b-j)\Gamma(1+a-b+j)}{\Gamma(b)\Gamma(a-2b+j+1)}\sum_{s=0}^{j}(-)^s\binom{j}{s}\frac{\Gamma(\frac{1}{2}a-b+\frac{1}{2}j+\frac{1}{2}s+\frac{1}{2})}{\Gamma(\frac{1}{2}a-\frac{1}{2}j+\frac{1}{2}s+\frac{1}{2})} \tag{3.5}$$

and

$$ {}_2F_1\left[\begin{array}{c} a,b \\ 1+a-b-j \end{array};-1\right] = 2^{-2b-j}\frac{\Gamma(1+a-b-j)}{\Gamma(a-2b-j+1)}\sum_{s=0}^{j}\binom{j}{s}\frac{\Gamma(\frac{1}{2}a-b-\frac{1}{2}j+\frac{1}{2}s+\frac{1}{2})}{\Gamma(\frac{1}{2}a-\frac{1}{2}j+\frac{1}{2}s+\frac{1}{2})} \tag{3.6} $$

for $j = 0, 1, 2, \ldots$. If we successively put $a = -n$, $b = -n-\nu$ and $a = -n-\nu$, $b = -n$ in (3.5) we obtain

$$ {}_2F_1\left[\begin{array}{c} -n,-n-\nu \\ 1+\nu+j \end{array};-1\right] $$

$$ = 2^{2n+2\nu+j}\frac{(-)^j\Gamma(1+n+\nu)\Gamma(1+\nu+j)}{\Gamma(1+n+\nu+j)\Gamma(n+2\nu+j+1)}\sum_{s=0}^{j}(-)^s\binom{j}{s}\frac{\Gamma(\frac{1}{2}n+\nu+\frac{1}{2}j+\frac{1}{2}s+\frac{1}{2})}{\Gamma(-\frac{1}{2}n-\frac{1}{2}j+\frac{1}{2}s+\frac{1}{2})} \tag{3.7} $$

and

$$ {}_2F_1\left[\begin{array}{c} -n,-n-\nu \\ 1-\nu+j \end{array};-1\right] = \frac{2^{2n+j}(-)^j n!}{j!(1+j)_n(1-\nu+j)_n}\sum_{s=0}^{j}(-)^s\binom{j}{s}\frac{\Gamma(\frac{1}{2}n-\frac{1}{2}\nu+\frac{1}{2}j+\frac{1}{2}s+\frac{1}{2})}{\Gamma(-\frac{1}{2}n-\frac{1}{2}\nu-\frac{1}{2}j+\frac{1}{2}s+\frac{1}{2})} \tag{3.8} $$

for $j = 0, 1, 2, \ldots$. Similarly, if we successively put $a = -n$, $b = -n-\nu$ and $a = -n-\nu$, $b = -n$ in (3.6) we obtain

$$ {}_2F_1\left[\begin{array}{c} -n,-n-\nu \\ 1+\nu-j \end{array};-1\right] = 2^{2n+2\nu-j}\frac{\Gamma(1+\nu-j)}{\Gamma(n+2\nu-j+1)}\sum_{s=0}^{j}\binom{j}{s}\frac{\Gamma(\frac{1}{2}n+\nu-\frac{1}{2}j+\frac{1}{2}s+\frac{1}{2})}{\Gamma(-\frac{1}{2}n-\frac{1}{2}j+\frac{1}{2}s+\frac{1}{2})} \tag{3.9} $$

and

$$ {}_2F_1\left[\begin{array}{c} -n,-n-\nu \\ 1-\nu-j \end{array};-1\right] = \frac{2^{2n-j}}{(1-\nu-j)_n}\sum_{s=0}^{j}\binom{j}{s}\frac{\Gamma(\frac{1}{2}n-\frac{1}{2}\nu-\frac{1}{2}j+\frac{1}{2}s+\frac{1}{2})}{\Gamma(-\frac{1}{2}n-\frac{1}{2}\nu-\frac{1}{2}j+\frac{1}{2}s+\frac{1}{2})} \tag{3.10} $$

for $j = 0, 1, 2, \ldots$.

3.2 *Evaluation of* $S_m(\pm\nu, \pm p)$

We consider in detail only the case corresponding to $S_m(\nu, p)$, the other cases being similar. We substitute (3.7) into (3.4) with $j = p + r$ and use the identity $(a+n)_r = (a+r)_n(a)_r/(a)_n$ to find

$$ S_m(\nu,p) = (-)^p 2^{2\nu+p}\sum_{r=0}^{m}\binom{m}{r}\frac{(-2x)^r\Gamma(1+\nu+r)}{(f)_r(1+\nu+p)_r\Gamma(1+2\nu+p+r)}\sum_{s=0}^{p+r}(-)^s\binom{p+r}{s} $$

$$ \times\sum_{n=0}^{\infty}\frac{(-x)^n}{n!}\frac{2^{2n}(1+\nu+r)_n}{(1+\nu+p+r)_n(1+2\nu+p+r)_n}\frac{\Gamma(\frac{1}{2}n+\nu+\frac{1}{2}p+\frac{1}{2}r+\frac{1}{2}s+\frac{1}{2})}{\Gamma(-\frac{1}{2}n-\frac{1}{2}p-\frac{1}{2}r+\frac{1}{2}s+\frac{1}{2})}. $$

Separating even and odd powers of x and making use of the identities

$$ (a)_{2n} = 2^{2n}(\tfrac{1}{2}a)_n(\tfrac{1}{2}a+\tfrac{1}{2})_n, \qquad (a)_{2n+1} = 2^{2n}a(\tfrac{1}{2}a+\tfrac{1}{2})_n(\tfrac{1}{2}a+1)_n, $$

$$ \frac{1}{\Gamma(-n+a)} = \frac{(-)^n(1-a)_n}{\Gamma(a)}, $$

we find after some routine algebra the desired result[1]

$$S_m(\nu,p) = (-)^p 2^{2\nu+p} \sum_{r=0}^{m} \binom{m}{r} \frac{(-2x)^r \Gamma(1+\nu+r)}{(f)_r (1+\nu+p)_r \Gamma(1+2\nu+p+r)} \sum_{s=0}^{p+r} (-)^s \binom{p+r}{s}$$
$$\times \left\{ A_r(\nu,p) F_r^{(1)} - \frac{4x(1+\nu+r) B_r(\nu,p)}{(1+\nu+p+r)(1+2\nu+p+r)} F_r^{(2)} \right\} \tag{3.11}$$

for $p = 0, 1, 2, \ldots$, where

$$A_r(\nu,p) = \frac{\Gamma(\nu+\frac{1}{2}p+\frac{1}{2}r+\frac{1}{2}s+\frac{1}{2})}{\Gamma(\frac{1}{2}s-\frac{1}{2}|p|-\frac{1}{2}r+\frac{1}{2})}, \qquad B_r(\nu,p) = \frac{\Gamma(\nu+\frac{1}{2}p+\frac{1}{2}r+\frac{1}{2}s+1)}{\Gamma(\frac{1}{2}s-\frac{1}{2}|p|-\frac{1}{2}r)}$$

and

$$F_r^{(1)} = {}_4F_5 \left[\begin{matrix} \frac{1}{2}+\frac{1}{2}\nu+\frac{1}{2}r,\ 1+\frac{1}{2}\nu+\frac{1}{2}r,\ \frac{1}{2}+\nu+\frac{1}{2}p+\frac{1}{2}r+\frac{1}{2}s,\ \frac{1}{2}+\frac{1}{2}p+\frac{1}{2}r-\frac{1}{2}s \\ \frac{1}{2},\ \frac{1}{2}+\frac{1}{2}\nu+\frac{1}{2}p+\frac{1}{2}r,\ 1+\frac{1}{2}\nu+\frac{1}{2}p+\frac{1}{2}r,\ \frac{1}{2}+\nu+\frac{1}{2}p+\frac{1}{2}r,\ 1+\nu+\frac{1}{2}p+\frac{1}{2}r \end{matrix} ; -x^2 \right],$$

$$F_r^{(2)} = {}_4F_5 \left[\begin{matrix} 1+\frac{1}{2}\nu+\frac{1}{2}r,\ \frac{3}{2}+\frac{1}{2}\nu+\frac{1}{2}r,\ 1+\nu+\frac{1}{2}p+\frac{1}{2}r+\frac{1}{2}s,\ 1+\frac{1}{2}p+\frac{1}{2}r-\frac{1}{2}s \\ \frac{3}{2},\ 1+\frac{1}{2}\nu+\frac{1}{2}p+\frac{1}{2}r,\ \frac{3}{2}+\frac{1}{2}\nu+\frac{1}{2}p+\frac{1}{2}r,\ 1+\nu+\frac{1}{2}p+\frac{1}{2}r,\ \frac{3}{2}+\nu+\frac{1}{2}p+\frac{1}{2}r \end{matrix} ; -x^2 \right].$$

An analogous procedure applied to $S_m(\nu,-p)$ produces

$$S_m(\nu,-p) = 2^{2\nu-p} \Gamma(1+\nu-p) \sum_{r=0}^{m} \binom{m}{r} \frac{(2x)^r (1+\nu)_r}{(f)_r \Gamma(1+2\nu-p+r)} \sum_{s-0}^{p-r} \binom{p-r}{s}$$
$$\times \left\{ A_r(\nu,-p) F_r^{(3)} - \frac{4x(1+\nu+r) B_r(\nu,-p)}{(1+\nu)(1+2\nu-p+r)} F_r^{(4)} \right\} \quad (p \geq m) \tag{3.12}$$

for positive integer p satisfying $p \geq m$, where

$$F_r^{(3)} = {}_4F_5 \left[\begin{matrix} \frac{1}{2}+\frac{1}{2}\nu+\frac{1}{2}r,\ 1+\frac{1}{2}\nu+\frac{1}{2}r,\ \frac{1}{2}+\nu-\frac{1}{2}p+\frac{1}{2}r+\frac{1}{2}s,\ \frac{1}{2}+\frac{1}{2}p-\frac{1}{2}r-\frac{1}{2}s \\ \frac{1}{2},\ \frac{1}{2}+\frac{1}{2}\nu,\ 1+\frac{1}{2}\nu,\ \frac{1}{2}+\nu-\frac{1}{2}p+\frac{1}{2}r,\ 1+\nu-\frac{1}{2}p+\frac{1}{2}r \end{matrix} ; -x^2 \right],$$

$$F_r^{(4)} = {}_4F_5 \left[\begin{matrix} 1+\frac{1}{2}\nu+\frac{1}{2}r,\ \frac{3}{2}+\frac{1}{2}\nu+\frac{1}{2}r,\ 1+\nu-\frac{1}{2}p+\frac{1}{2}r+\frac{1}{2}s,\ 1+\frac{1}{2}p-\frac{1}{2}r-\frac{1}{2}s \\ \frac{3}{2},\ 1+\frac{1}{2}\nu,\ \frac{3}{2}+\frac{1}{2}\nu,\ 1+\nu-\frac{1}{2}p+\frac{1}{2}r,\ \frac{3}{2}+\nu-\frac{1}{2}p+\frac{1}{2}r \end{matrix} ; -x^2 \right].$$

We observe that (3.12) is valid provided $p \geq m$. This is a consequence of (3.9) holding for non-negative j, which when j is set equal to $p-r$ with $0 \leq r \leq m$, yields the requirement $p \geq m$. If $m > p$, then we must use (3.9) for $0 \leq r \leq p$ and (3.7) for $p+1 \leq r \leq m$; see the remark at the end of this section.

Use of (3.8) and (3.10) in the same manner yields the representations for $S_m(-\nu,\pm p)$ given by

$$S_m(-\nu,p) = \frac{(-2)^p}{p!} \sum_{r=0}^{m} \binom{m}{r} \frac{(-2x)^r (1+\nu)_r}{(f)_r (1-\nu+p)_r (1+p)_r} \sum_{s-0}^{p+r} (-)^s \binom{p+r}{s}$$
$$\times \left\{ C_r(-\nu,p) F_r^{(5)} - \frac{4x(1+\nu+r) D_r(-\nu,p)}{(1+\nu)(1-\nu+p+r)(1+p+r)} F_r^{(6)} \right\} \tag{3.13}$$

[1]We omit the subscript s on the quantities $A_r(\nu,p)$, $B_r(\nu,p)$ and $F_r^{(j)}$ to ease the notation.

and

$$S_m(-\nu,-p)=2^{-p}\sum_{r=0}^{m}\binom{m}{r}\frac{(2x)^r(1+\nu)_r}{(f)_r(1-\nu-p)_r}\sum_{s-0}^{p-r}\binom{p-r}{s}$$

$$\times\left\{C_{-r}(-\nu,-p)F_r^{(7)}-\frac{4x(1+\nu+r)D_{-r}(-\nu,-p)}{(1+\nu)(1-\nu-p+r)}F_r^{(8)}\right\}\quad(p\geq m),\tag{3.14}$$

where

$$C_r(\nu,p)=\frac{\Gamma(\frac12\nu+\frac12p+\frac12|r|+\frac12s+\frac12)}{\Gamma(\frac12\nu-\frac12|p|-\frac12r+\frac12s+\frac12)},\quad D_r(\nu,p)=\frac{\Gamma(\frac12\nu+\frac12p+\frac12|r|+\frac12s+1)}{\Gamma(\frac12\nu-\frac12|p|-\frac12r+\frac12s)}.$$

Here we have defined the functions $F_r^{(j)}$ $(5\leq j\leq 8)$ by

$$F_r^{(5)}={}_5F_6\left[\begin{matrix}1,\ \frac12+\frac12\nu+\frac12r,\ 1+\frac12\nu+\frac12r,\ \frac12-\frac12\nu+\frac12p+\frac12r+\frac12s,\ \frac12+\frac12\nu+\frac12p+\frac12r-\frac12s\\ \frac12+\frac12\nu,\ 1+\frac12\nu,\ \frac12+\frac12p+\frac12r,\ 1+\frac12p+\frac12r,\ \frac12-\frac12\nu+\frac12p+\frac12r,\ 1-\frac12\nu+\frac12p+\frac12r\end{matrix};-x^2\right],$$

$$F_r^{(6)}={}_5F_6\left[\begin{matrix}1,\ 1+\frac12\nu+\frac12r,\ \frac32+\frac12\nu+\frac12r,\ 1-\frac12\nu+\frac12p+\frac12r+\frac12s,\ 1+\frac12\nu+\frac12p+\frac12r-\frac12s\\ 1+\frac12\nu,\ \frac32+\frac12\nu,1+\frac12p+\frac12r,\ \frac32+\frac12p+\frac12r,\ 1-\frac12\nu+\frac12p+\frac12r,\ \frac32-\frac12\nu+\frac12p+\frac12r\end{matrix};-x^2\right],$$

$$F_r^{(7)}={}_4F_5\left[\begin{matrix}\frac12+\frac12\nu+\frac12r,\ 1+\frac12\nu+\frac12r,\ \frac12-\frac12\nu-\frac12p+\frac12r+\frac12s,\ \frac12+\frac12\nu+\frac12p-\frac12r-\frac12s\\ \frac12,\ \frac12+\frac12\nu,\ 1+\frac12\nu,\ \frac12-\frac12\nu-\frac12p+\frac12r,\ 1-\frac12\nu-\frac12p+\frac12r\end{matrix};-x^2\right],$$

$$F_r^{(8)}={}_4F_5\left[\begin{matrix}1+\frac12\nu+\frac12r,\ \frac32+\frac12\nu+\frac12r,\ 1-\frac12\nu-\frac12p+\frac12r+\frac12s,\ 1+\frac12\nu+\frac12p-\frac12r-\frac12s\\ \frac32,\ 1+\frac12\nu,\ \frac32+\frac12\nu,\ 1-\frac12\nu-\frac12p+\frac12r,\ \frac32-\frac12\nu-\frac12p+\frac12r\end{matrix};-x^2\right].$$

We note that $C_{-r}(-\nu,-p)=1$ and $D_{-r}(-\nu,-p)=\frac12(s-p+r-\nu)$.

As in (3.12), we note that the representation (3.14) is valid for $p\geq m$. To deal with the situation corresponding to $m>p$ we must write the sums $S_m(\pm\nu,-p)$ as

$$S_m(\pm\nu,-p)=\sum_{r=0}^{p}\binom{m}{r}\frac{x^r(1+\nu)_r}{(f)_r(1\pm\nu-p)_r}\sum_{n=0}^{\infty}\frac{(-x)^n}{n!}\frac{(1+\nu+r)_n}{(1+\nu)_n}\,{}_2F_1\left[\begin{matrix}-n,-n-\nu\\ 1\pm\nu-(p-r)\end{matrix};-1\right]$$

$$+\sum_{r=p+1}^{m}\binom{m}{r}\frac{x^r(1+\nu)_r}{(f)_r(1\pm\nu-p)_r}\sum_{n=0}^{\infty}\frac{(-x)^n}{n!}\frac{(1+\nu+r)_n}{(1+\nu)_n}\,{}_2F_1\left[\begin{matrix}-n,-n-\nu\\ 1\pm\nu+(r-p)\end{matrix};-1\right].$$

Then in the first series we employ the ${}_2F_1$ summations in (3.7) and (3.9) for $p-r\geq 0$, and in the second series we employ the summations in (3.8) and (3.10) for $r-p>0$. We omit these details.

4. Concluding remarks

When $m=0$ the representations (3.11)–(3.14) reduce to the corresponding cases given in (2.2)–(2.5). In the cases $r=0$ and $r=1$ the hypergeometric functions appearing in $S_m(-\nu,p)$ and $S_m(\pm\nu,-p)$ contract to yield lower-order functions; this is not the case for $S_m(\nu,p)$ when $p\geq 2$.

For example, in the case of $S_m(-\nu,p)$ the ${}_5F_6$ functions contract to ${}_3F_4$ functions when $r=0$ and to ${}_4F_5$ functions when $r=1$.

In the case $p=0$, $m=1$, for example, we find that the ${}_4F_5$ functions contract to simpler ${}_0F_1$ functions to yield the result

$$\begin{aligned} S_1(\nu,0) &= e^{-x}\sum_{n=0}^{\infty}\frac{x^n L_n^{(\nu)}(x)}{(1+\nu)_n}\,\frac{(f+1)_n}{(f)_n} \\ &= \left(1+\frac{x}{f}\right){}_0F_1\left[\begin{array}{c}\text{—}\\ 1+\nu\end{array};-x^2\right]-\frac{x^2}{(1+\nu)f}\,{}_0F_1\left[\begin{array}{c}\text{—}\\ 2+\nu\end{array};-x^2\right] \\ &= \Gamma(1+\nu)x^{-\nu}\left\{\left(1+\frac{x}{f}\right)J_\nu(2x)-\frac{x}{f}\,J_{\nu+1}(2x)\right\}, \end{aligned}$$

where we have employed the duplication formula for the gamma function and expreesed ${}_0F_1$ in terms of the J-Bessel function..

Finally, we mention that the summations $S_m(\pm\nu,\pm p)$ in (3.11)–(3.14) have been verified numerically with the help of *Mathematica*.

References

[1] Yu. A. Brychkov, Two series containing the Laguerre polynomials, Integral Transforms and Special Functions (2013) http://dx.doi.org/10.1080/1065 2469.2013.777532.

[2] H. Exton, Summation formulae involving Laguerre polynomials, J. Comput. Appl. Math. **100** (1998) 225–227.

[3] Y. S. Kim, A. K. Rathie and R. B. Paris, On a new class of summation formulae involving the Laguerre polynomial, Integral Transforms and Special Functions **23(6)** (2012) 435–444.

[4] Y. S. Kim, A. K. Rathie and R. B. Paris, A note on a series containing Laguerre polynomials, Miskolc Math. Notes (2014) [to appear].

[5] A. P. Prudnikov, Yu. A. Brychkov and O. I. Marichev, Integrals and Series: Special Functions, Vol. 3, Gordon and Breach, New York, 1988.

[6] M. A. Rakha and A. K. Rathie, Generalizations of classical summation theorems for the series ${}_2F_1$ and ${}_3F_2$ with applications, Integral Transforms and Special Functions **22(11)** (2011) 823–840.

[7] A. K. Rathie and R. B. Paris, Extension of some classical summation theorems for the generalized hypergeometric series with integral parameter differences, J. Classical Anal. **3(2)** (2013) 109–127.

[8] L. J. Slater, Generalized Hypergeometric Functions, Cambridge University Press, Cambridge 1966.